

Saccharomyces - B



FUNCTIONAL
NUTRIENTS®

Clinical Applications

- Supports Gastrointestinal-based Immunity*
- Supports Integrity and Function of Mucosal Cells*
- Supports Healthy Gut Flora*

*Saccharomyces - B is a lactose-free, stomach acid-resistant, stable, European patent-pending formula containing DNA-verified Saccharomyces boulardii. This probiotic yeast supports other probiotic organisms in addition to intestinal barrier function and integrity. Research also suggests that Saccharomyces boulardii supports normal immune responses.**

All Personalized Medical Solutions Formulas Meet or Exceed cGMP Quality Standards

Discussion

Saccharomyces - B is manufactured using a process that involves controlled temperature and low vacuum, eliminating the shock of freeze-drying. The drying procedure, which does not modify the water content of the preparation, slows down aging, cell deterioration, and contamination, leading to increased stability and greater resistance to gastric acid. This process does not require or contain dairy products or lactose.*

Saccharomyces boulardii (*S boulardii*) is a natural, non-pathogenic yeast that may help support and maintain the healthy ecology of the small and large intestines. In a 2010 systematic review and meta-analysis of 31 randomized placebo-controlled treatment arms in 27 trials comprising 5029 adults, "*S boulardii* was found to be significantly efficacious and safe in 84% of those treatment arms."^[1] A study also suggested that *S boulardii* can be used safely and effectively in children three months and older.^[2] A double-blind, randomized, placebo controlled study of children ages 3-59 months produced statistically significant positive results suggesting that *S boulardii* may play an important role in supporting normal gastrointestinal function in young children.*^[3]

Extensively researched and published in European and American peer-reviewed journals, *Saccharomyces boulardii* appears to function through several mechanisms of action. With regard to maintaining normal GI function and transit time, research suggests that *S boulardii* secretes a protease that may assist in directly degrading bacterial toxins and stimulating antibody production against those toxins.^[4-6] *S boulardii* is also believed to support the natural inflammatory response, exert a trophic effect on intestinal mucosa, and positively support the immune system.^[3] These actions further support and help maintain the normal health and function of the intestinal brush border.*

This probiotic yeast appears to support normal gastrointestinal flora and integrity^[7,8]; promote production of intestinal enzymes and secretory IgA^[9]; and support cytokine balance through its effects on IL-8, IL-6, IL-10, NF-kappaB, TNF-alpha, and PPAR-gamma.^[6,10,11] Research suggests a role for *S boulardii* in the production of health-promoting short-chain fatty acids, including butyrate.*^[12] Research suggests that butyrate, in turn, plays an important role in maintaining the integrity, function, and normal flora of the intestines.^[13]

Following a multi-dose study in healthy volunteers, the concentration of *S boulardii* in feces increased rapidly, reaching a steady state by day three in all subjects. Following completion of administration, the *S boulardii* population declined consistently and was cleared from the bowel within five to seven days, indicating that the probiotic yeast did not permanently colonize the gastrointestinal tract.^[14] Saccharomyces - B provides 10 billion colony-forming units of *S boulardii* in a two-capsule dose.*

***These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.**



Supplement Facts

Serving Size: 2 Capsules
Servings Per Container: 30

	Amount Per Serving	%Daily Value
<i>Saccharomyces boulardii</i> (10 billion CFU†)	500 mg	**

** Daily Value not established.

Other Ingredients: HPMC (capsule), microcrystalline cellulose, stearic acid, magnesium stearate, silica, and medium-chain triglyceride oil.

Europe Patent Pending #05425084.0

† Colony-Forming Units

Directions

Take one to two capsules per day, or as directed by your healthcare practitioner.

Consult your healthcare practitioner prior to use. Not for use by infants under three months of age. Do not use if you have severe immune suppression or are taking antifungal medication. Do not use if tamper seal is damaged.

References

1. McFarland LV. Systematic review and meta-analysis of *Saccharomyces boulardii* in adult patients. *World J Gastroenterol*. 2010 May 14;16(18):2202-22. [PMID: 20458757]
2. Vandenas Y, Brunser O, Szajewska H. *Saccharomyces boulardii* in childhood. *Eur J Pediatr*. 2009 Mar;168(3):253-65. Review. [PMID: 19096876]
3. Riaz M, Alam S, Malik A, et al. Efficacy and Safety of *Saccharomyces boulardii* in Acute Childhood Diarrhea: A Double Blind Randomised Controlled Trial. *Indian J Pediatr*. 2011 Oct 14. [PMID: 21997865]
4. Can M, Beşirbellioğlu BA, Avci İY, et al. Prophylactic *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhea: a prospective study. *Med Sci Monit*. 2006 Apr;12(4):P119-22. [PMID: 16572062]
5. Castagliuolo I, Riegler MF, Valenick L, et al. *Saccharomyces boulardii* protease inhibits the effects of *Clostridium difficile* toxins A and B in human colonic mucosa. *Infect Immun*. 1999 Jan;67(1):302-7. [PMID: 9864230]
6. Im E, Pothoulakis C. [Recent advances in *Saccharomyces boulardii* research]. *Gastroenterol Clin Biol*. 2010 Sep;34 Suppl 1:S62-70. Review. French. [PMID: 20889007]
7. Krasowska A, Murzyn A, Dyjankiewicz A, et al. The antagonistic effect of *Saccharomyces boulardii* on *Candida albicans* filamentation, adhesion and biofilm formation. *FEMS Yeast Res*. 2009 Dec;9(8):1312-21. [PMID: 19732158]
8. *Saccharomyces boulardii*. Professional Monograph. Natural Standard Database <http://naturalstandard.com/databases/herbssupplements/sboulardii.asp>. Accessed February 14, 2012.
9. Buts JP, Bernasconi P, Vaerman JP, et al. Stimulation of secretory IgA and secretory component of immunoglobulins in small intestine of rats treated with *Saccharomyces boulardii*. *Dig Dis Sci*. 1990 Feb;35(2):251-6. [PMID: 2302983]
10. Dahan S, Dalmaso G, Imbert V, et al. *Saccharomyces boulardii* interferes with enterohemorrhagic *Escherichia coli*-induced signaling pathways in T84 cells. *Infect Immun*. 2003 Feb;71(2):766-73. [PMID: 12540556]
11. Pothoulakis C. Review article: anti-inflammatory mechanisms of action of *Saccharomyces boulardii*. *Aliment Pharmacol Ther*. 2009 Oct 15;30(8):826-33. Review. [PMID: 19706150]
12. Schneider SM, Girard-Pipau F, Filippi J, et al. Effects of *Saccharomyces boulardii* on fecal short-chain fatty acids and microflora in patients on long-term total enteral nutrition. *World J Gastroenterol*. 2005 Oct 21;11(39):6165-9. [PMID: 16273644]
13. Lewis K, Lutgendorff F, Phan V, et al. Enhanced translocation of bacteria across metabolically stressed epithelia is reduced by butyrate. *Inflamm Bowel Dis*. 2010 Jul;16(7):1138-48. [PMID: 20024905]
14. Blehaut H, Massot J, Elmer GW, et al. Disposition kinetics of *Saccharomyces boulardii* in man and rat. *Biopharm Drug Dispos*. 1989 Jul-Aug;10(4):353-64. [PMID: 2758101]

Does Not Contain

Wheat, gluten, corn, soy, animal or dairy products, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically-modified organisms (GMOs), artificial colors, artificial sweeteners, or preservatives.

***These statements have not been evaluated by the Food and Drug Administration.
This product is not intended to diagnose, treat, cure, or prevent any disease.**

Personalized Medical Solutions
22525 SE 64th Place Ste 200 Issaquah, WA 98027
Report any adverse reactions to 425-391-3376

DRS-109
REV. 03/30/18